


FORM PTO-1390		U.S. Department of Commerce Patent and Trademark Office	Attorney's Docket No. 2577-112
<b>TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371</b>			U.S. Application No. (if known, see 37 CFR 1.5) <b>10/019992</b>
INTERNATIONAL APPLICATION NO. PCT/SG99/00061	INTERNATIONAL FILING DATE June 22, 1999	PRIORITY DATE CLAIMED	
<b>TITLE OF INVENTION</b> Detection of <i>Salmonella Enteritidis</i>			
<b>APPLICANT(S) FOR DO/EO/US</b> Hwei-Sing KWANG, Wei LIU, Su-Shing Sharon LOW, Kwang Yeng Hilda LOH			
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:			
1. <input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371 2. <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). 4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) a. <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). b. <input checked="" type="checkbox"/> has been transmitted by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US) 6. <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> have been transmitted by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has <b>NOT</b> expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).			
<b>ITEMS 11. TO 16. below concern other document(s) or information included:</b>			
11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input checked="" type="checkbox"/> A <b>FIRST</b> preliminary amendment. <input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment. 14. <input type="checkbox"/> A substitute specification. 15. <input type="checkbox"/> A change of power of attorney and/or address letter. 16. <input checked="" type="checkbox"/> Other items or information: Courtesy copy of International Application No. PCT/SG99/00061 w/ attached International Search Report.			

U.S. APPLICATION NO. (if known, see 37 CFR 1.50) <b>10/019992</b>		INTERNATIONAL APPLICATION NO PCT/SG99/00061		ATTORNEY DOCKET NO 2577-112											
17. <input checked="" type="checkbox"/> The following fees are submitted: <b>Basic National Fee (37 CFR 1.492(a)(1)-(5)):</b> Search Report has been prepared by the EPO or JPO \$ 890.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) \$ 710.00 No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)) \$ 740.00 Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$ 1,040.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4) \$ 100.00				<table border="1" style="width:100%; border-collapse: collapse;"> <tr> <th style="width:50%;">CALCULATIONS</th> <th style="width:50%;">PTO USE ONLY</th> </tr> <tr> <td> </td> <td> </td> </tr> <tr> <td> </td> <td> </td> </tr> <tr> <td> </td> <td> </td> </tr> <tr> <td> </td> <td> </td> </tr> </table>		CALCULATIONS	PTO USE ONLY								
CALCULATIONS	PTO USE ONLY														
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$ 890.00											
Surcharge of \$130.00 for furnishing the oath or declaration later than [ ] 20 [ ] 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$											
Claims	Number Filed	Number Extra	Rate												
Total Claims	19 -20 =	0	X \$18.00	\$											
Independent Claims	3 -3 =	0	X \$84.00	\$											
Multiple dependent claim(s) (if applicable)			+ \$280.00	\$											
TOTAL OF ABOVE CALCULATIONS =				\$890.00											
Reduction by 1/2 for filing by small entity, if applicable. Applicant claims small entity status.				\$445.00											
SUBTOTAL =				\$445.00											
Processing fee of \$130.00 for furnishing the English translation later [ ] 20 [ ] 30 than months from the earliest claimed priority date (37 CFR 1.492(f)).				\$											
TOTAL NATIONAL FEE =				\$445.00											
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property				\$											
TOTAL FEES ENCLOSED =				\$445.00											
				Amount to be refunded	\$										
				charged	\$										
a. <input type="checkbox"/> A check in the amount of \$ _____ to cover the above fees is enclosed. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. 02-2135 in the amount of \$445.00 to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 02-2135. A duplicate copy of this sheet is enclosed. d. <input type="checkbox"/> Payment by credit card. (Form PTO-2038 enclosed.) <b>NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.</b>															
SEND ALL CORRESPONDENCE TO: CUSTOMER No 6449 Barbara G. Ernst Rothwell, Figg, Ernst & Manbeck 555 13th St., N.W. Washington, D.C. 20004 Phone: 202/783-6040				<div style="text-align: center;">         Signature        Jeffrey L. Ihnen        Name        28,957        Registration Number     </div>											

<b>IN THE UNITED STATES PATENT AND TRADEMARK OFFICE</b>	Application Number	§371 of PCT/SG99/00061
	Filing Date	June 22, 1999
	First Named Inventor	Hwei-Sing KWANG
	Group Art Unit	Unassigned
	Examiner Name	Unassigned
	Attorney Docket Number	2577-112
Title of the Invention: DETECTION OF SALMONELLA ENTERITIDIS		

**PRELIMINARY AMENDMENT**

Assistant Commissioner for Patents  
 Washington, D.C. 20231

Dear Sir:


Please amend the above-identified U.S. patent application as follows:

IN THE CLAIMS:

Please cancel claims 20-34.

**REMARKS**

The above is being made to make amendment to the claims prior to examination on the merits. The amendment does not add to or depart from the original disclosure, or constitute prohibited new matter.

RESPECTFULLY SUBMITTED,					
NAME AND REG. NUMBER	Jeffrey L. Ihnen, Registration No. 28,957				
SIGNATURE				DATE	December 26, 2001
Address	Rothwell, Figg, Ernst & Manbeck Suite 701-East, 555 13th Street, N.W.				
City	Washington	State	D.C.	Zip Code	20004
Country	U.S.A.	Telephone	202-783-6040	Fax	202-783-6031

Nucleotide sequence of *sefA*

ATTTTGTAATATGCGTAAATCAGCATCTGCAGTAGCAGTTCTTGCTTTAATTGCAT  
GTGGCAGTGCCACGCAGCTGGCTTTGTTGGTAACAAAGCAGAGGTTTCAGGCAGC  
GGTACTATTGCAGCTCAGAATACAACATCAGCCAACTGGAGTCAGGATCCTGGC  
TTTACAGGGCCTGCTGTTGCTGCTGGTCAGAAAGTTGGTACTCTCAGCATTACTG  
CTACTGGTCCACATAACTCAGTATCTATTGCAGGTAAAGGGGCTTCGGTATCTGG  
TGGTGTAGCCACTGTCCCGTTTCGTTGATGGACAAGGACAGCCTGTTTTCCGTGGG  
CGTATTCAGGGAGCCAATATTAATGACCAAGCAAATACTGGAATTGACGGGCTTG  
CAGGTTGGCGAGTTGCCAGCTCTCAAGAAACGCTAAATGTCCCTGTCACAACCTT  
TGGTAAATCGACCCTGCCAGCAGGTACTTTCAGTGCAGACCTTCTACGTTTCAGCAG  
TATCAAAAC

(SEQ ID NO:1)

WO 00/78995

PCT/SG99/00061

2

Amino acid sequence for SEF14:

1

MRKSASAVAVLALACGSAHAAGFVGKAEVQAAVTIAAQNTTSANW  
SQDPGFTGPAVAAGQKVGTLSTATGPHNSVSIAGKGASVSGGVATVP  
FVDGQGQPVFRGRIQGANINDQANTGIDGLAGWRVASSQETLNVPVT  
TFGKSTLPAGTFTATFYVQQYQN

165

Amino acid sequence for the C128 fragment of SEF14:

AAQNTTSANWSQDPGFTGPAVAAGQKVGTLSTATGPHNSVSIAGKGA  
SVSGGVATVPFVDGQGQPVFRGRIQGANINDQANTGIDGLAGWRVASS  
QETLNVPVTTFGKSTLPAGTFTATFYVQQYQN  
(SEQ ID NO:3)

Amino Acid Sequence of *S. enteritidis* Flagellin Antigen

LTQNNLNKSQSSLSSAIERLSSGLRINSAKDDAAGQAIANRFTS  
NIKGLTQASRNANDGISIAQTTEGALNEINNNLQRVRELSVQATNGTNSDSLKSIQD  
EIQQRLEEIDRVSNQTQFNGVKVLSQDNQMKIQVGANDGETITIDLQKIDVKSLGLDG  
FNVNGPKEATVGDLSKSFKNVTGYDTYAAGADKYRVDINSGAVVTDAAAPDKVYVNAA  
NGQLTTDDAENNTAVDLFKTTKSTAGTAEAKAIRGAIKGGKEGDTFDYKGVFTTIDTK  
TGDDGNGKVSTTINGEKVTLTVADIATGATDVNAATLQSSKNVYTSVVNGQFTFDDKT  
KNESAKLSGLEANNAVKGESKITVNGAEYTANATGDKITLAGKTMFIDKTASGVSTLI  
NEDAAAANKSTANPLASIDSALSKVDAVRSSLGAIQNRFDSAITNLGNTVTNLNSARS  
RIEDADYATEVSNMSKAQILQQAGTSVLAQANQVPQNVLSLLR

(SEQ ID NO:4)

90 amino acid fragment of *S. enteritidis* flagellin antigen

TAEAKAIRGAIKGGKEGDTFDYKGVFTTIDTKTGDDGNGKVSTTINGEKVTLTVADIA  
TGATDVNAATLQSSKNVYTSVVNGQFTFDDKT

(SEQ ID NO:5)

fragment A: 69 amino acids (aa 258-327 of SEQ ID NO:4)

KEGDTFDYKGVFTTIDTKTGDDGNGKVSTTINGEKVTLTVADIATGATDVNAATLQSSKN  
VYTSVVNGQ

(SEQ ID NO:6)

fragment B: 40 amino acids (aa 276-316 of SEQ ID NO:4)

KTGDDGNGKVSTTINGEKVTLTVADIATGATDVNAATLQS

(SEQ ID NO:7)

fragment C: 27 amino acids (aa 279-306 of SEQ ID NO:4)

DGNGKVSTTINGEKVTLTVADIATGAT

(SEQ ID NO:8)

fragment D: 11 amino acids (aa 285-296 of SEQ ID NO:4)

STTINGEKVTL

(SEQ ID NO:9)

DETECTION OF SALMONELLA ENTERITIDISFIELD OF THE INVENTION

This invention relates to a method for detecting *Salmonella enteritidis* in poultry and in their eggs. More specifically, the invention is directed to a method for detecting *S. enteritidis* which comprises contacting a biological sample obtained from poultry suspected of containing *S. enteritidis* with a fragment of the *S. enteritidis* fimbrial protein or a fragment of the *S. enteritidis* flagellin protein which specifically recognizes *S. enteritidis* antibodies present in the sample and discriminates between *S. enteritidis* and other *Salmonella spp.*

BACKGROUND OF THE INVENTION

*Salmonella enteritidis*, an agent which causes salmonellosis in poultry, can be transmitted vertically from laying hens to eggs. Consumption of eggs or meat contaminated with the organism can lead to food poisoning in humans. This is a worldwide problem in public health; in the U.S. alone, more than a million cases of salmonellosis are reported annually. Outbreaks in the elderly and in young children can be especially dangerous, resulting in severe gastroenteritis and possibly fatal septicemia.

Possible *Salmonella* virulence factors include fimbrial structures, which are gene products involved in the invasion of eukaryotic cells, and lipopolysaccharides. Another factor is the flagella, which confer motility to the bacterium and so contribute to the bacterium's colonization.

In view of the large number of cases of *Salmonella enteritidis* cases reported each year, there is an obvious need for a reliable method for detecting laying flocks infected with *S. enteritidis*. Bacteriological techniques for the isolation of *S. enteritidis*, such as those disclosed by Williams, J.E., and A.D. Whittemore, *Avian Disease*, 20:728 (1976), are laborious, time-consuming and costly. False-negative results can arise when *S. enteritidis* is overgrown by other *Salmonella* serotypes present in the samples. In addition, these methods may not identify all birds infected with *S. enteritidis* because *Salmonella* excretion is intermittent.

There are some existing serological methods, such as the serum plate test (SPT), latex agglutination test (LAT) and enzyme linked immunosorbent assay (ELISA), which are rapid and easy to perform, and antibodies against *S. enteritidis* have been found at relatively high levels in the sera of infected chickens, making antibody detection practical. Most ELISAs utilize either lipopolysaccharides (LPS) or flagella antigens for the detection of antibody against *S. enteritidis*. The use of these antigens, however, has resulted in false positive results in recent reports, making discrimination between *Salmonella* serotypes difficult.



See Christopher, J., et al. *J. Clin. Microbiol.* 34:792-797 (1996); van Zijderveld, Fred, *J. Clin. Microbiol.* 36:2560-2566 (1992); Barrow, P.A., *Epidemiology and Infection* 109:361-369 (1992); and Barrow, P.A., *Int. J. Food Microbiol.* 21:55-68 (1994). Accordingly, further methods are sought.

#### SUMMARY OF THE INVENTION

This invention is directed to a method for detecting *S. enteritidis* which comprises contacting a biological sample obtained from poultry suspected of being infected with *S. enteritidis* with an antigenic fragment of an *S. enteritidis* fimbrial protein or an antigenic fragment of an *S. enteritidis* flagellin protein, which fragment specifically recognizes antibodies against *S. enteritidis* present in the sample and discriminates between antibodies against *S. enteritidis* and antibodies against other *Salmonella* spp, under conditions sufficient for the formation of an immunological complex between *S. enteritidis* antibodies present in the sample and the antigenic fragment and then detecting the formation of such a complex.

Another embodiment of this invention comprises a method for detecting *S. enteritidis* which comprises contacting a biological sample obtained from poultry suspected of being infected with *S. enteritidis* with a combination of an antigenic fragment of an *S. enteritidis* fimbrial protein and an antigenic fragment of an *S. enteritidis* flagellin protein, each of which fragments specifically recognizes antibodies against *S.*

*enteritidis* present in the sample and discriminates between antibodies against *S. enteritidis* and antibodies against other *Salmonella spp*, under conditions sufficient for the formation of an immunological complex between *S. enteritidis* antibodies present in the sample and either or both of the antigenic fragments and then detecting the formation of such a complex.

This invention further is directed to a diagnostic kit, which comprises an antigenic fragment of the *S. enteritidis* fimbrial protein, an antigenic fragment of the *S. enteritidis* flagella protein, or both such fragments, wherein said fragment(s), when combined with a biological sample obtained from poultry suspected of being infected with *S. enteritidis*, specifically recognizes antibodies against *S. enteritidis* present in said sample and discriminates between antibodies against *S. enteritidis* and antibodies against other *Salmonella spp*.

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 sets forth the DNA sequence encoding the *S. enteritidis* fimbrial protein SEF14.

Figure 2A sets forth the amino acid sequence of the *S. enteritidis* fimbrial protein SEF14 and Figure 2B sets forth the amino acid sequence of the fragment identified herein as C128.

Figure 3 is a chart illustrating eight subfragments of fimbrial protein SEF14 that were tested for reactivity to *S. enteritidis*.

Figure 4 is a representation of the full-length *S. enteritidis* flagellin antigen and the fragments obtained from the full-length sequence as described in Example 2.

Figure 5 sets forth the amino acid sequence of the full length *S. enteritidis* flagellin protein illustrated in Figure 4, as well as the amino acid sequence encoded by the 270 bp fragment and the amino acid sequence of each of four subfragments which were isolated as described in Example 2.

#### DETAILED DESCRIPTION OF THE INVENTION

##### Determination of antigenic fragments

###### A. Antigenic fragments of fimbrial antigen

A fimbrial antigen designated SEF14 was first described in 1994 (Thorns, C.J., et al., *J. Clin. Microbiol.* 28:2409-2414). The SEF14-encoding gene, designated *sefA*, was shown to be limited in distribution to serotypes belonging to *Salmonella* serogroup D. Expression of the SEF14 antigen as a surface structure has been detected only in *S. enteritidis*, *S. dublin*, *S. blegdam* and *S. moscow*, but *S. enteritidis* is the only serotype that can be isolated from poultry, and SEF14 fimbriae is expressed by all *S. enteritidis* strains. It is known that antibody against SEF14 is developed following infections with *S. enteritidis*. The entire sequence of the *sefA* gene is known and has been published in the literature (Thorns, C.J., et al., *J. Clin. Microbiol.* 4(34):792-797 (1996)) and in GenBank (accession number L03833). The DNA sequence of the *sefA* gene is provided

in Figure 1 and is identified herein as SEQ ID NO. 1. The amino acid sequence for SEF14 is provided in Figure 2A. This sequence is identified herein as SEQ ID NO. 2.

To determine which portions of SEF14 specifically recognize *S. enteritidis*, the *sefA* gene was amplified from genomic DNA with primers designated from Genbank sequence data. The resulting amplified fragment was cloned into an expression vector. The vector was chosen such that the *sefA* gene product would be expressed as a fusion protein. A preferred fusion partner is glutathione -S- transferase of *Schistosoma japonica* (GST).

The reactivity of the resultant recombinant GST-SEF14 fusion protein was tested with sera from chickens experimentally infected with *S. enteritidis* and other *Salmonella* serotypes in immunoblot. GST-SEF14 was recognized by all *S. enteritidis*-infected sera, and weak reaction bands also were obtained with sera from infection of *S. senftenberg*, *S. amsterdam*, *S. java*, *S. pullorum*, *S. typhimurium*, *S. agona* and *S. oranienberg*.

No reactions were found with *S. haardt*, *S. montevideo*, *S. paratyphimurium*, *S. emek* and pre-infection sera. From these data it appeared that the SEF14 fusion protein shares some common epitopes with other *Salmonella* serotypes which caused these cross-reactions.

In order to locate a partial SEF14 fragment specific to *S. enteritidis*, eight sub-fragments of the 165 amino acid SEF14, as shown in Figure 3, were expressed as fusion proteins with GST. The reactivity

of each fragment was tested by immunoblot assays. Of the three fragments containing the N-terminus of SEF14, only the full-length SEF14, named F165 (165 aa), was recognized by all sera from *S. enteritidis* infections.

The other two fragments, N150 (150 aa) and N136 (136 aa), were not recognized by some of the *S. enteritidis* sera, suggesting that the failure probably was due to the deletion of some dominant epitopes in the C-terminus between aa 151 and aa 165.

Of the four fragments containing the C-terminus of SEF14, two fragments, C145 (145 aa; aa 121-165) and C128 (128 aa; aa 38-165), reacted with all *S. enteritidis* sera. The other two fragments, C95 (95 aa) and C88 (88 aa), were not detected by some of the *S. enteritidis* sera, which suggested that the epitopes between aa 37 and aa 70 are associated with antibody binding. The amino acid sequence of the C128 fragment is provided in Figure 2B and is identified herein as SEQ ID NO. 3.

The hydrophilicity and antigenicity of the SEF14 amino acid residues were analyzed by the Hydrophilicity Plot : Kyte-Doolittle and Antigenic Index : Jameson-Wolf computer based programs. These programs assist in the prediction of the characteristics of the protein from the knowledge of the sequence. The regions of amino acids 41-53, 144-153 and 159-165 are hydrophilic, and it is believed that they represent antigenic epitopes. The deletion of these regions from SEF14 results in a polypeptide with decreased antigenicity.

The specificity of the SEF sub-fragments also was tested with sera from chickens experimentally infected with various *Salmonella* serotypes. Sub-fragment C128 did not react with sera from any *Salmonella* serotype other than *S. enteritidis*. Subfragment C145 reacted weakly with sera from infection of *S. senftenberg*, *S. java*, *S. pullorum*, *S. amsterdam* and *S. paratyphimurium*. The deletion of N-terminus aa 1-36 of F165 made subfragment C128 more specific than F165 or C145. When the F165 fragment was tested, cross-reactions could be observed, and slight cross-reactions also were observed when the C145 fragment was tested. In contrast, when C128 was tested, no cross reaction was seen. These results indicate that the region bounded by aa 1-36 contains at least one epitope shared by different *Salmonella* serotypes. These studies are described in detail in Example 1, below. From the foregoing, therefore, a desirable fragment within the scope of this invention is one which consists essentially of an antigenic subfragment of the C145 fragment of SEF14 which specifically recognizes *S. enteritidis* antibodies in a biological sample obtained from poultry and distinguishes between *S. enteritidis* and other *Salmonella* species. Such subfragments include the C128 subfragment, which consists essentially of amino acids 37-165 of the SEF14 amino acid sequence, and further subfragments thereof which specifically recognize *S. enteritidis* antibodies in a biological sample obtained from poultry and distinguish between *S. enteritidis* and other *Salmonella* species. Also included within the scope of this invention are polypeptides which

The specificity and sensitivity of subfragment C128 also was studied using ELISA with sera obtained from guinea pigs hyperimmunized with *Salmonella spp.* and *Enterobacteriaceae spp.* and sera from chickens experimentally infected with *Salmonella spp.* With sera from guinea pigs, the OD value for *S. enteritidis* was at least 2 times as high as those of other sera with the exception of *S. dublin*, which had a similar OD value due to cross-reaction of SEF14 fimbriae by *S. dublin*. This cross-reactivity with *S. dublin* is not a problem in terms of being able to use C128 as a diagnostic for the identification of poultry infected with *S. enteritidis* because *S. dublin* does not infect, and so cannot be isolated from, chickens. With sera obtained from chickens, all sera from chickens infected with *S. enteritidis* showed an OD value at least two times higher than those obtained from sera from chickens infected with other *Salmonella* serotypes. These data indicate that the reactivity of C128 in ELISA was the same as the reactivity in immunoblot. The cut-off OD value for positive reaction was three times that of the average OD of normal sera.

The C128 subfragment of F165 thus is a useful tool for the detection of *S. enteritidis* in poultry.

B. Antigenic fragment of flagellin antigen

The amino acid sequence of the flagellin region of *S. enteritidis* (nucleotides 754-1024 of *S. enteritidis* strain designated 13076 and deposited with and available from the ATCC as accession number U12963) is shown in Figure 5, identified as SEQ ID NO:4. A DNA fragment encoding the flagellin region was amplified and cloned into an expression vector. The vector was chosen such that the desired flagellin gene product would be expressed as a fusion protein. A preferred fusion partner is glutathione -S- transferase of *Schistosoma japonica* (GST). The resultant GST-flagellin protein recombinant protein was expressed. The amino acid sequence of the 90 amino acid fragment of the flagellin protein is provided in Figure 5. The sequence is identified herein as SEQ ID NO:5.

This recombinant protein was recognized by *Salmonella dublin* as well as by *S. enteritidis*, but not by any of the other *Salmonella* spp. tested.

To determine if a smaller sequence could be used to specifically identify *S. enteritidis* in poultry samples, the sequence of the flagellin portion of the recombinant protein was compared to the sequences of the flagellin domains of other *Salmonella* spp. and the specific regions within the protein that showed the most variation from the other sequences were identified. Four subfragments of the flagellin domain were expressed as fusion proteins ranging in size from 27.2kD to 31.6 kD, and the reactivity of each fragment



was tested. Each fragment was shown to specifically detect and discriminate samples from poultry infected with *S. enteritidis* from uninfected samples. These four subfragments are shown in Figure 5 and identified herein as SEQ ID NOS: 6-9.

The 90 amino acid fragment and each of the four subfragments illustrated in Figure 5 thus are useful in the method of the present invention. Also useful is an antigenic fragment of any of these sequences or an amino acid sequence which corresponds to one of these sequences which comprises a conservative amino acid substitution for at least one amino acid in the sequence, provided that with said substitution(s) the sequence specifically recognizes *S. enteritidis* antibodies in a biological sample obtained from poultry and distinguishes between *S. enteritidis* and other *Salmonella* species.

#### Detection of *S. enteritidis* infections

The antigenic fragments of *S. enteritidis* fimbrial and flagellin proteins can be used to detect *S. enteritidis* infections in samples obtained from poultry. Preferred samples are sera samples or yolk samples from poultry eggs. A sample is contacted with an antigenic fragment of *S. enteritidis* fimbrial or flagellin proteins in accordance with the present invention for a time and under conditions sufficient for the formation of an immunological complex between *S. enteritidis* antibodies present in the sample and the antigenic fragment. The formation of the resulting complex can be assayed either by direct detection

Using direct detection methods, the antigen-antibody complexes can be assayed, for example by ELISA, in accordance with standard techniques. The direct detection assays can use labeled antigenic fragment or labeled anti-*S. enteritidis* antibodies. Labeling of the antibody or antigenic fragment can be conducted using standard labeling techniques. The detectable label can be a fluorescent compound, a radioactive element, an enzyme capable of producing a detectable reaction product, or gold. The selected antigenic fragment of *S. enteritidis* fimbrial or flagellin proteins can be labeled, for example, with a radioactive isotope. The sample then is contacted with a radioactively labeled antigen fragment of *S. enteritidis* fimbrial or flagellin proteins in accordance with the present invention for a time and under conditions sufficient for the formation of an immunological complex between *S. enteritidis* antibodies present in the sample and the antigenic fragment, such that antibody-antigen complexes which form are labeled with the radioactive label. The formed complex then can be assayed by immunoblotting. Alternatively, *S.*

*enteritidis* antibodies present in the sample can be labeled with a directly detectable label, such as a fluorescent compound or gold, or can be conjugated to an enzyme commonly used for colorimetric or fluorescent detection, such as alkaline phosphatase. Unlabeled antigenic fragment of *S. enteritidis* fimbrial or flagellin proteins then can be coated onto the microtiter plates and contacted with the biological sample containing labeled *S. enteritidis* antibodies so that, again, antibody-antigen complexes which form are labeled with the detectable label. Detection of the formed antigen-antibody complexes that are labeled can be conducted, for example, by a standard ELISA assay or other sensitive detection system.

Using indirect detection methods, the antigen-antibody complexes can be assayed, for example, by ELISA or immunoblotting in accordance with standard techniques. Antigenic fragments of *S. enteritidis* fimbriae or flagellin proteins can be coated onto microtiter plates which can provide a solid phase for capturing anti-*S. enteritidis* antibodies present in the sample. Alternatively, antigenic fragments can be provided in solution for immunoblotting. The fragment is contacted with the biological sample under conditions sufficient for the formation of an immunological complex between *S. enteritidis* antibodies present in the sample and the antigenic fragment. Antibody-antigen complexes that form then can be detected using secondary antisera conjugated to detectable labels, such as fluorescent-labeled antibodies, enzyme-conjugated antibodies, including

horseradish peroxidase-conjugated antibodies or alkaline phosphatase-conjugated antibodies, radioactive tracer-labeled antibodies or gold-labeled antibodies.

In an alternative to these assay techniques, monoclonal antibodies to the antigenic fragments of *S. enteritidis* of the present invention can be generated in accordance with standard techniques. The monoclonal antibodies then can be used to coat the wells of microtiter plates which then are contacted with a biological sample suspected of containing *S. enteritidis* and antigens in the sample will bind to the monoclonal antibodies and can be detected using a detectable label as described above.

A preferred assay for use in the present invention is an ELISA as described above. A second preferred assay is a lateral flow format assay, which can easily be provided in the form of a rapid test kit. Such kits, which are commercially available for other purposes, allow the absorption of fluids via application to a pre-determined well containing a strip of antigen, or as in this case, an antigenic fragment, on a membrane and then inserting the well into a device which has a window above the antigen-coated membrane. The sample is added to the well such that it flows across the membrane, allowing any antibodies present in the sample to interact with the antigen in the membrane. A secondary antibody which is labeled with a detectable label, such as gold, also is incorporated into the membrane and, upon wetting of the membrane with the sample, is mobilized and binds to any primary antibody present in the sample. Any positive result

obtained is observed by the appearance of a band on the membrane produced by the precipitation of the gold onto the membrane which is visualized through the window. The lateral flow format assay also can be modified by coating monoclonal antibodies generated against the desired antigenic fragment onto the membrane.

Western blotting, dot blots and quartz crystal technology also can be utilized for the purpose of detection of the antigen or monoclonal antibody.

Assays using the C128 fimbrial fragment and assays using one the flagellin fragments can be used independently for the detection of *S. enteritidis* infections in poultry and their eggs. It also can be useful to use the fimbriae and flagellin fragments in combination. Detectable antibodies against SEF14 indicate an early stage infection, while antibodies to the flagellin proteins typically are detected later in the infection, as the antibody to flagella is generated later but has a longer existence. Thus, by testing a sample with antigenic fragments of each of the fimbriae protein and flagellin protein, one can detect the presence of an infection from its beginning stages to its mature stages and thus ensure that no infections go undetected.

#### Diagnostic Kits

The invention further comprises diagnostic kits which can be used to detect and discriminate *S. enteritidis* infections in samples from poultry. The kits comprise an antigenic fragment of the *S. enteritidis* fimbrial protein or flagellin domain, as

described above, or can comprise both an antigenic fragment of the *S. enteritidis* fimbriae protein and an antigenic fragment of the *S. enteritidis* flagellin domain. The kit further can comprise a label. Suitable labels include labels which can be attached to the antigenic fragment or to antibodies present in the biological sample to be tested, such as an enzyme, gold, fluorescent compound, or radioactive element. Alternatively, the label can be provided in the form of a labeled secondary antibody for use in an indirect detection method, such as enzyme-conjugated antibodies, including horseradish peroxidase conjugated antibodies or alkaline phosphatase labeled antibodies, gold-labeled antibodies, fluorescently-labeled antibodies, or radiotracer-labeled antibodies.

In one preferred embodiment, the kit comprises the essential elements for a lateral flow format assay as described above.

The invention is further described in the following examples, which are provided for illustrative purposes and are not intended to be construed as limiting.

#### Example 1

##### Bacterial strains

*S. enteritidis* strains 2/93 phage type 4, 119/95 phage type 4, 330/96 phage type 11a and 131/97 phage type 37, 40/97 phage type 1, and 296/96 phage type 9b were provided by the Primary Production Department,

Singapore, and isolate 94/6510 was provided by the U.S. Department of Agriculture. All of these strains are publicly available. These strains were used for the experimental infection of chicken.

#### Serum samples

##### Sera Group 1

Guinea pigs were inoculated with different strains of bacterium to obtain serum specific to the following: *S. enteritidis* (group D), *S. dublin* (group C), *S. kentucky* (group C3), *S. heidelberg* (group B), *S. newport* (group C3), *S. typhimurium copenhagen* (group B), *S. cholersasius*, *S. anatum* (group E1), *S. cerro* (group K), *Escherichia coli*, *Aeromonas hydrophilus*, *Pasteurella multocida*, *Klebsiella pneumonia*, *Proteus mirabilis*, *Yersinia enterocoli*, *Citrofreundii*, *shigella sonnei*, *Serratia marcescens* and uninfected guinea pig serum.

##### Sera group 2

Chickens were infected with various serotypes of *Salmonella* obtained from the Singapore General Hospital: *S. dublin* (group D), *S. typhimurium* (group B), *S. anatum* (group E1), *S. pullorum* (group D), *S. java* (group B), *S. paratyphi A* (group A), *S. haardt* (group C3), *S. hadar* (group C2), and *E. coli*.

##### Sera group 3

The six *S. enteritidis* strains listed above were prepared as an overnight broth which was diluted to  $1 \times 10^8$  colony forming units (cfu). This broth in turn was diluted further, resulting in  $10^7$  and  $10^5$  concentrations of the six cultures. Twelve ten-week old chickens were

divided into six groups of two, and one chicken in each group was inoculated with 1 ml of the  $10^5$  bacterial broth and the second in each group inoculated with 1 ml of the  $10^7$  bacterial broth. Sera from each chicken was obtained at seven day intervals and the flock was monitored for *Salmonella* shedding from pre-inoculation to two weeks post-inoculation.

#### Sera group 4

Forty samples of *S. enteritidis* negative sera were obtained from specific pathogen-free chickens ranging in age from 1 day old chicks to ten week old chickens.

Specifically, the forty samples comprised sera obtained from ten one day-old chicks, fourteen four week old chicks, and sixteen ten week old chicks.

#### Sera group 5

Twenty five serum samples were collected from chickens in the field by the Primary Production Department (PPD), Singapore. These samples were collected from farm chickens in Malaysia in which *S. enteritidis* infections had been identified.

#### Recombinant *S. enteritidis* antigen

The nucleotide sequence of the *sefA* gene is available through Genbank (accession number L03833). Oligonucleotide primers for the C128 fragment of the fimbrial SEF14 antigen were designed and synthesized as follows. Primers were designed with a Bam H1 restriction site in the 5' end for the forward primer and an EcoR1 restriction site at the 3' end for the reverse primer.

forward primer:



5' TGC AGC TCA GAA TAC AAC ATC A 3' (starting from base 112)

reverse primer:

5' GTT TTG ATA CTG CTG AAC GTA (end at base 495)

The forward and reverse primers are identified herein as SEQ ID NOS: 10 and 11, respectively.

The Beckman OLIGO 1000M DNA synthesizer was used in the synthesis. Using genomic DNA extracted from *S. enteritidis* strain 13076 ATCC, DNA fragments were amplified by PCR. The amplified DNA was cloned into pGEX-4T-3 expression vector (available from Pharmacia Biotech, Uppsala, Sweden). Clones containing inserts were sequenced to ensure the correctness of the reading frame. Proteins fused to GST were expressed in *E. coli* strain JM105, obtained from Amersham Pharmacia Biotech, Uppsala, Sweden; catalog no. 27-1550-01. For the purification of the fusion protein, the bacterial cell pellet was subjected to GST affinity column purification (Pharmacia Biotech), following the instructions provided by the manufacturer. To attain higher purity, the protein was loaded onto SDS-PAGE gel and the protein band visualized by a 1 minute stain with Coomassie brilliant blue and de-stained in de-ionized water. The band was excised and eluted in de-ionized water. The protein obtained through such gel purification was used as antigen in an ELISA test.

#### Immunoblot

Western blot analysis was carried out according to conventional format with some modifications. Purified fusion protein was mixed with sample buffer containing

0.1 mol/L DTT in final concentration. The mixture was denatured at 100°C for 3 minutes and separated by SDS-PAGE. The separated protein on gel was transferred to a nitrocellulose membrane by overnight electrophoresis under 20 volts. The membrane was blocked with 5% skim milk and stripped.

Serum samples were diluted at 1:400, added to each of the strips individually, and incubated at room temperature for 1 hour, followed by the incubation of rabbit anti-chicken IgG peroxidase conjugate for one hour and then by color development in 3'3'-diaminobenzidine tetrahydrochloride (DAB) substrate. All fimbrial fragments were found to be positive in reactivity.

#### ELISA

An ELISA was preformed using a conventional format. Purified protein was coated on 96-well flat bottom plates (NUNC) in carbonate buffer, pH 9.6 at 50 ng/100 µl per well. After blocking with 1% BSA, serum samples, diluted 1:200, were added and incubated at 37°C for 15 minutes, followed by incubation of secondary IgG peroxidase conjugate at 37°C for 15 minutes and finally by addition of substrate OPD. Results were expressed as the optical density (OD) at 492 nm by ELISA reader (Bio-dot).

The C128 antigen fragment was mapped to be specific and sensitive to *S. enteritidis* infection. the cutoff value for a positive reaction was 3 times the OD value of that of uninfected chicken sera. The fragment was tested against Group 2, where only *S.*

### ELISA procedure

Immulon microtiter plates (available from Dynatech Laboratories Inc. Chantilly, Va.) were coated with 50 ng/100 microliters/well recombinant protein in 0.1M sodium bicarbonate buffer (ph 9.6). The plates were incubated at 37°C for 4 hours and then refrigerated until further use. The plates were washed 4 times with ELISA washing solution (phosphate buffered saline, 0.05% Tween 20[PBST]) and excess binding sites were saturated with 1% bovine serum albumin (BSA) fraction V (Sigma) in phosphate buffered saline (pH 7.4) for 1 hour at 37°C. After four washings, 100 microliters of the test serum sample(1:200 dilution) in 1% BSA-PBS buffer were added to each well, and the plates were incubated for 10 minutes at 37°C. Following subsequent washing of the wells, 100 microliters of anti-chicken immunoglobulins conjugated with horseradish peroxidase (KPL) were added to each well and the plates were incubated at 37°C for 10 minutes. The wells were washed again and 100 microliters of the substrate solution (2'2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid (ABTS) with H<sub>2</sub>O<sub>2</sub>) were added. The color reaction was allowed to proceed at room temperature for 10 minutes, and the absorbance of each well at 405 nm (ABTS) respectively was recorded in an automatic ELISA plate reader.

#### Mapping of the *S. enteritidis* partial protein

In order to assess if the obtained partial protein, shown in Figure 5 as SEQ. ID NO:5, was the smallest fragment usable for the purpose of detecting the presence of *S. enteritidis* in poultry or their

eggs, primers were designed to amplify specific regions within the partial fragment which showed the greatest sequence variation between *S. enteritidis* and other flagellin domains of *Salmonella* serotypes identified in Asten et al., *J. Bacteriology* 177(6):1610 (1995). Primers (set forth in the table below) and the resulting fragments were cloned using the same method set forth above in vector pGEX-2T. The four resulting partial protein fragments were designated FLP-A, FLP-B, FLP-C and FLP-D. These partial protein fragments also were purified by affinity chromatography and gel excision. These partial protein fragments were quantified and subsequently coated as 50 ng per well. Characterization of these partial protein fragments was conducted with the use of chicken sera in ELISA. Serum obtained from uninfected chickens yielded negative results, while serum obtained from chicken infected with *S. enteritidis* yielded positive results.

Table 1 Nucleotide sequences of primers used

<i>Fragments</i>	<i>Primers Used</i>	<i>Nucleotides amplified</i>
A: Forward A: Reverse	5'AAGGATCCAAAGGTGGTAAGGAAGGA5' 5'CCGAATTCTTTGTCACCGTTCACTAC3'	775bp to 982bp'
B: Forward B: Reverse	5'AAGGATCCAAAAGTGGTGATGACGGT3' 5'CCGAATTCGCTTGATTGTAAGGTAGC3'	828bp TO 948bp
C: Forward C: Reverse	5'AAGGATCCGACGGTAATGGTAAGGTT3' 5'CCGAATTCGTCGCGCCAGTGGCAAT3'	838bp to 919 bp

The primers listed above are identified herein as SEQ ID NOS: 12-19, respectively.

The average reading was calculated for the negative sera used in these assays and the standard deviation calculated. A range of standard deviations were calculated, where the average of the negative sera was added to one, two, three, four or five standard deviations. It was found that the average of the negative reference sera plus two of their standard deviations was sufficient in value to include all negatives and all the known positive sera values obtained were above that value. The cut off value (detection limit) of the ELISA was defined as the mean value of the negative reference sera plus two times their standard deviation, which is able to discriminate the positive sera and also field infected sera.

Differences in the sets of data collected were statistically analyzed by the t-test (Sigma Plot program). The value obtained indicates whether the differences seen in two sets are significantly different. P values of less than 0.01 are considered to be statistically different.

Characterization and specificity of the *S. enteritidis* flagellin fusion protein

Gel purified GST-*S. enteritidis* antigen was immunologically characterized by Western Blot. The fusion protein was tested with sera raised from guinea pigs from Group 1 sera. Of the ten *Salmonella* serotypes tested, only *S. enteritidis* and *S. dublin* of the same serogroup displayed reactivity. The sera raised from other serotypes and other organisms in Group 1 showed no reactivity. All sera were screened at a dilution of 1:50. The fusion protein then was screened against experimentally infected and non-infected sera obtained from chickens. Group 3 sera all reacted positively and Group 4 sera showed no reaction.

Development of ELISA for *S. enteritidis* using GST-*S. enteritidis* fusion protein

Gel purified GST-*S. enteritidis* flagellin antigen fragment fusion protein was diluted in ELISA coating buffer and used as coating antigen on microtiter plates at 50 ng per well overnight. The coated antigen was characterized by the guinea pig sera from Group 1. Subsequent characterization was carried out using chicken sera. Sera from *S. enteritidis*-infected chickens had the highest absorbance value, which was calculated to have a positive/negative ratio of 4.59 times greater than the average of the negative references.

Non-infected chicken sera from Group 3 (sera obtained as pre-bleeds from the chickens prior to infection) then were tested against the coated antigen.

All yielded almost basal readings. Infected flock sera from Group 3 which were of 5 different phage types yielded 100% positive results. Thus, this test is able to detect *S. enteritidis* of different phage types.

Sera from Group 2, obtained from the flock experimentally infected with *S. haardt*, *S. typhimurium*, *S. hadar*, *S. typhi* and *S. java*, gave negative results with the test antigen. These data confirm that there are no cross-reactions with other *Salmonella* serotypes.

#### Comparison with commercially available IDEXX *Salmonella enteritidis* test kit

Group 5 sera were evaluated using both the IDEXX commercial test kit (IDEXX Laboratories Inc., Westbrook, ME) and the antigen of the present invention. Using the IDEXX kit, 3/28 sera were detected; using the antigen of the present invention, 14/28 samples were identified as serologically positive. The antigen of the present invention thus showed a detection rate which was 78.6% greater than obtained using the commercially available test kit.

#### ELISA results: mapped fragments

The *S. enteritidis* fusion protein was further mapped to assess the smallest region which could still detect and discriminate infected from non-infected sera. A total of four fragments, shown in Figure 5 as fragments A-D and having molecular weights (inclusive



of GST) 31.6kD, 30.4 kD, 29 kD, and 27.2 kD, were cloned and screened with the selected positive and negative sera. Eight serum samples from chicken experimentally infected with *S. enteritidis* and eight samples of SPF (specific pathogen free) sera were used to qualify each fragment for its sensitivity. Positive/negative (P/N) ratios for the fragments were determined. The average P/N ratio for each of the fragments was as follows:

fragment A: 3.23  
 fragment B: 2.95  
 fragment C: 3.02  
 fragment D: 2.97

These figures reflect a similar reactivity pattern to that of the original fusion antigen, which has a P/N ratio of 3.01.

The fragments were tested for specificity with other *Salmonella* serotypes present in the samples of serum in Group 2. None of the fragments reacted with any of the serotypes other than *S. enteritidis*. The results are shown in Table 2 below:

Table 2  
 Fragment Reactivity with Serotype Specific Serum

Fragments/ Serum	270	A	B	C	D
<i>S. enteritidis</i>	+	+	+	+	+
<i>S. dublin</i>	-	-	-	-	-
<i>S. typhimurium</i>	-	-	-	-	-

<i>S.anatum</i>	-	-	-	-	-
<i>S.pullorum</i>	-	-	-	-	-
<i>S.java</i>	-	-	-	-	-
<i>S.paratyphi</i>	-	-	-	-	-
<i>S.haardt</i>	-	-	-	-	-
<i>S.hadar</i>	-	-	-	-	-
<i>E.coli</i>	-	-	-	-	-

## Claims:

1. A method for detecting *Salmonella enteritidis* in a biological sample obtained from poultry which comprises

contacting said biological sample with an antigenic fragment of *S. enteritidis* fimbrial protein or an antigenic fragment of *S. enteritidis* flagellin protein under conditions sufficient for the formation of an immunological complex between *S. enteritidis* antibodies present in said sample and said fragment, and

detecting the formation of such a complex; wherein said fragment specifically recognizes *S. enteritidis* antibodies present in the sample and discriminates between *S. enteritidis* and other *Salmonella* spp.

2. The method of claim 1, wherein said sample comprises sera or egg yolk.

3. The method of claim 1, wherein said sample is contacted with a fragment of *S. enteritidis* fimbrial protein.

4. The method of claim 3, wherein said fragment is provided as a fusion polypeptide wherein an additional polypeptide is fused to said fragment.

9. The method of claim 7, wherein said fragment consists essentially of the amino acid sequence of SEQ ID NO:5, an antigenic fragment thereof, or a sequence which corresponds to said sequence and comprises a conservative amino acid substitution for at least one amino acid in said sequence.

10. The method of claim 7, wherein said fragment consists essentially of the amino acid sequence of SEQ ID NO:6, an antigenic fragment thereof or a sequence which corresponds to said sequence which comprises a conservative amino acid substitution for at least one amino acid in said sequence.

11. The method of claim 7, wherein said fragment consists essentially of the amino acid sequence of SEQ ID NO:7, an antigenic fragment thereof or a sequence which corresponds to said sequence which comprises a conservative amino acid substitution for at least one amino acid in said sequence.

12. The method of claim 7, wherein said fragment consists essentially of the amino acid sequence of SEQ ID NO:8, an antigenic fragment thereof or a sequence which corresponds to said sequence which comprises a conservative amino acid substitution for at least one amino acid in said sequence.

13. The method of claim 7, wherein said fragment consists essentially of the amino acid sequence of SEQ ID NO:9, an antigenic fragment thereof or a sequence which corresponds to said sequence which comprises a conservative amino acid substitution for at least one amino acid in said sequence.

14. The method of claim 1, wherein said fragment is labeled with a detectable label.



of SEQ ID NO:7, an antigenic fragment thereof or a sequence which corresponds to said sequence which comprises a conservative amino acid substitution for at least one amino acid in said sequence.

21. An isolated fragment of *S. enteritidis* flagellin protein consisting of the amino acid sequence of SEQ ID NO:8, an antigenic fragment thereof, or a sequence which corresponds to said sequence and comprises a conservative amino acid substitution for at least one amino acid in said sequence.

22. An isolated fragment of *S. enteritidis* flagellin protein consisting of the amino acid sequence of SEQ ID NO:9, an antigenic fragment thereof or a sequence which corresponds to said sequence and comprises a conservative amino acid substitution for at least one amino acid in said sequence.

23. A kit comprising (a) a fragment of *S. enteritidis* fimbrial or flagellin protein which specifically recognizes *S. enteritidis* antibodies present in a biological sample obtained from poultry suspected of being infected with *S. enteritidis* and discriminates between antibodies from *S. enteritidis* and other *Salmonella* spp. and (b) a detectable label.

24. A kit comprising a fragment of *S. enteritidis* fimbrial protein and a fragment of *S. enteritidis* flagellin protein, each of which fragments specifically recognizes *S. enteritidis* antibodies present in a

biological sample obtained from poultry suspected of being infected with *S. enteritidis* and discriminates between antibodies from *S. enteritidis* and other *Salmonella* spp..

25. A kit in accordance with claim 24, which further comprises a detectable label.

26. A method for detecting *Salmonella enteritidis* in a biological sample obtained from poultry which comprises

contacting said biological sample with an antigenic fragment of *S. enteritidis* fimbrial protein and an antigenic fragment of *S. enteritidis* flagellin protein under conditions sufficient for the formation of an immunological complex between *S. enteritidis* antibodies present in said sample and either or both of said fragments, and

detecting the formation of such a complex or complexes;  
wherein each of said fragments specifically recognizes *S. enteritidis* antibodies present in the sample and discriminates between *S. enteritidis* and other *Salmonella* spp.

27. A method for detecting *Salmonella enteritidis* in a biological sample obtained from poultry which comprises

contacting a first portion of said biological sample with an antigenic fragment of *S. enteritidis* fimbrial protein under conditions sufficient for the



30. The method of claim 26 or 27, wherein said fragment of *S. enteritidis* fimbrial protein consists essentially of a subfragment of aa 121-165 of the amino

34. The method of claim 26 or 27, wherein said label comprises a fluorescent compound, a radioactive element, an enzyme capable of producing a reaction detectable compound or gold.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number  
WO 00/78995 A1

- (51) International Patent Classification<sup>7</sup>: C12Q 1/10, (74) Agent: ELLA CHEONG & G. MIRANDAH; P.O. Box 0931, Raffles City, Singapore 911732 (SG).  
C07K 14/255
- (21) International Application Number: PCT/SG99/00061 (81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 22 June 1999 (22.06.1999)
- (25) Filing Language: English
- (26) Publication Language: English
- (71) Applicant (*for all designated States except US*): INSTITUTE OF MOLECULAR AGROBIOLOGY [SG/SG]; The National University of Singapore, 1 Research Link, Singapore 117604 (SG).
- (72) Inventors; and (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (75) Inventors/Applicants (*for US only*): KWANG, Hwei-Sing [SG/SG]; 87 Pasir Panjang Hill, #02-09 Kent New Mark, Singapore 118892 (SG). LIU, Wei [SG/SG]; Block 508, Apartment 8265, West Coast Drive, Singapore (SG). LOW, Su-Shing, Sharon [SG/SG]; 17C Nassim Road, 01-04 Nassim Park, Singapore 258394 (SG). LOH, Kwang, Yeng, Hilda [SG/SG]; 73 Cavenagh Road, #08-372, Singapore 229624 (SG).
- Published:  
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: DETECTION OF *SALMONELLA ENTERITIDIS*

(57) Abstract: A method for detecting *Salmonella enteritidis* in poultry and their eggs comprises contacting a biological sample obtained from poultry suspected of containing *S. enteritidis* with a fragment of a *S. enteritidis* fimbrial protein or a fragment of a *S. enteritidis* flagellin protein which specifically recognizes *S. enteritidis* antibodies present in the sample and discriminates between *S. enteritidis* and other *Salmonella* spp.










PTO/PCT Rec'd 26 DEC 2001




WO 00/78995 A1

WO 00/78995

1/2

	N-terminus	C-terminus	
N136			136aa
N150			150aa
F165			165aa
C145			145aa
C128			128aa
C95			95aa
C88			88aa
M124+			124aa
M100			100aa

2/2



<b>DECLARATION AND POWER OF ATTORNEY FOR UTILITY OR DESIGN PATENT APPLICATION</b> (37 CFR 1.63)	Attorney Docket No.	2577-112
	First Named Inventor	Hwei-Sing Kwang .
	COMPLETE IF KNOWN	
	Application Number	10/018,892
	Filing Date	December 26, 2001
	Group Art Unit	
	Examiner Name	

☐ Declaration Submitted with Initial Filing
 ☒ Declaration Submitted after Initial Filing

As a below named inventor, I hereby declare that:

My residence, mailing address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: DETECTION OF SALMONELLA ENTERITIDIS the specification of which was filed on June 22, 1999 as PCT International Application Number PCT/SG99/00061.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56, including for continuation-in-part applications, material information which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Numbers	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
			<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)

I or we hereby appoint the registered practitioner(s) associated with Customer No. 6449 to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith. Direct all correspondence to Customer Number 6449.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF SOLE OR FIRST INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle [if any]) <u>1-00 Hwei-Sing</u>		Family Name or Surname <u>KWANG</u>	
Inventor's Signature <u>[Signature]</u>		Date <u>25/6/02</u>	
Residence: City <u>Singapore</u> <u>SGX</u>		Country <u>Singapore</u>	Citizenship <u>USA</u>
Mailing Address <u>54 West Coast Crescent, #03-02, West Bay Condimin.</u>			
Mailing Address			
City <u>Singapore</u>		Postal Code <u>428037</u>	Country <u>Singapore</u>
NAME OF SECOND INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle [if any]) <u>Wei</u>		Family Name or Surname <u>LIU</u>	
Inventor's Signature <u>[Signature]</u>		Date <u>25-06-2002</u>	
Residence: City <u>Singapore</u> <u>SGX</u>		Country <u>Singapore</u>	Citizenship <u>China</u>
Mailing Address <u>Block 508, West Coast Drive, #08-265 B1K280 Toh Guan Rd #12-205 Singapore 600280</u>			
Mailing Address			
City <u>Singapore</u>		Postal Code <u>120508</u>	Country <u>Singapore</u>
NAME OF THIRD INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle [if any]) <u>Su-Shing Sharon</u>		Family Name or Surname <u>LOW</u>	
Inventor's Signature		Date	
Residence: City <u>Singapore</u> <u>SGX</u>		Country <u>Singapore</u>	Citizenship <u>Singapore</u>
Mailing Address <u>17C Nassim Road, #01-04 Nassim Park</u>			
Mailing Address			
City <u>Singapore</u>		Postal Code <u>258394</u>	Country <u>Singapore</u>
NAME OF FOURTH INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle [if any]) <u>Hilda Kwanyeng</u>		Family Name or Surname <u>LOH</u>	
Inventor's Signature		Date	
Residence: City <u>Singapore</u> <u>SGX</u>		Country <u>Singapore</u>	Citizenship <u>Singapore</u>
Mailing Address <u>73, Cavanaugh Road, #08-372</u>			
Mailing Address			
City <u>Singapore</u>		Postal Code <u>229624</u>	Country <u>Singapore</u>

NAME OF SOLE OR FIRST INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any)) Hwei-Sing		Family Name or Surname KWANG	
Inventor's Signature		Date	
Residence: City Singapore		Country Singapore	Citizenship USA
Mailing Address 54 West Coast Crescent, #03-02, West Bay Condo Min.			
Mailing Address			
City Singapore		Postal Code 426037	Country Singapore
NAME OF SECOND INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any)) Wei		Family Name or Surname LIU	
Inventor's Signature		Date	
Residence: City Singapore		Country Singapore	Citizenship China
Mailing Address Block 508, West Coast Drive, #08-285			
Mailing Address			
City Singapore		Postal Code 120508	Country Singapore
NAME OF THIRD INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any)) Su-Shing Sharon		Family Name LOW or Surname	
Inventor's Signature		Date	
Residence: City Singapore		Country Singapore	Citizenship Singapore
Mailing Address 17C Nassim Road, #01-04 Nassim Park			
Mailing Address			
City Singapore		Postal Code 258394	Country Singapore
NAME OF FOURTH INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any)) Hilda Kwanyeng		Family Name LOH or Surname	
Inventor's Signature <i>H-LL</i>		Date 18-6-2002	
Residence: City Singapore		Country Singapore	Citizenship Singapore
Mailing Address 73, Cavanagh Road, #08-372			
Mailing Address			
City Singapore		Postal Code 229624	Country Singapore



<b>DECLARATION AND POWER OF ATTORNEY FOR UTILITY OR DESIGN PATENT APPLICATION</b> (37 CFR 1.63)		Attorney Docket No.	2577-112
		First Named Inventor	Hwei-Sing Kwang
		COMPLETE IF KNOWN	
		Application Number	10/018,892
		Filing Date	December 26, 2001
		Group Art Unit	
<input type="checkbox"/> Declaration Submitted with Initial Filing		<input checked="" type="checkbox"/> Declaration Submitted after Initial Filing	Examiner Name

As a below named inventor, I hereby declare that:

My residence, mailing address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: DETECTION OF SALMONELLA ENTERITIDIS the specification of which was filed on June 22, 1999 as PCT International Application Number PCT/SG99/00061.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56, including for continuation-in-part applications, material information which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or any PCT international application having a filing date before that of the application on which priority is claimed.


Prior Foreign Application Numbers	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
			<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>

I hereby claim the benefit under 36 U.S.C. 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)

I or we hereby appoint the registered practitioner(s) associated with Customer No. 6449 to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith. Direct all correspondence to Customer Number 6449.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

<b>NAME OF SOLE OR FIRST INVENTOR:</b>		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any)) Hwei-Sing		Family Name or Surname KWANG	
Inventor's Signature		Date	
Residence: City Singapore		Country Singapore	Citizenship USA
Mailing Address 54 West Coast Crescent, #03-02, West Bay Condo Min.			
Mailing Address			
City Singapore		Postal Code 428037	Country Singapore
<b>NAME OF SECOND INVENTOR:</b>		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any)) Wei		Family Name or Surname LIU	
Inventor's Signature		Date	
Residence: City Singapore		Country Singapore	Citizenship China
Mailing Address Block 508, West Coast Drive, #08-265			
Mailing Address			
City Singapore		Postal Code 120508	Country Singapore
<b>NAME OF THIRD INVENTOR:</b>		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any)) Su-Shing Sharon		Family Name LOW or Surname	
Inventor's Signature 		Date 9th July 2002	
Residence: City Singapore		Country Singapore	Citizenship Singapore
Mailing Address 17C Nassim Road, #01-04 Nassim Park			
Mailing Address			
City Singapore		Postal Code 258394	Country Singapore
<b>NAME OF FOURTH INVENTOR:</b>		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any)) Hilda Kwanyeng		Family Name LOH or Surname	
Inventor's Signature		Date	
Residence: City Singapore		Country Singapore	Citizenship Singapore
Mailing Address 73, Cavanaugh Road, #08-372			
Mailing Address			
City Singapore		Postal Code 229624	Country Singapore

PTO/PCT Rec'd 06 AUG 2002

FB

SEQUENCE LISTING

<110> Kwang, Hwei-Sing  
Liu, Wei  
Low, Su-Shing Sharon  
Loh, Hilda Kwanyeng

<120> Detection of Salmonella Enteritidis

<130> 2577-112

<150> PCT/SG99/00061

<151> 1999-06-22

<160> 19

<170> PatentIn version 3.1

<210> 1

<211> 505

<212> DNA

<213> Salmonella enteritidis

<400> 1

atattgtaaat atgcgtaaat cagcatctgc agtagcagtt cttgctttta ttgcatgtgg	60
cagtgccac gcagctggct ttgttggttaa caaagcagag gttcaggcag cggttactat	120
tgcagctcag aatacaacat cagccaactg gagtcaggat cctggcttta cagggcctgc	180
tggttgctgct ggtagcagaag ttgttactct cagcattact gctactggtc cacataactc	240
agtatctatt gcaggtaaag gggcttcggt atctgggtgg gttagccactg tcccgttcgt	300
tgatggacaa ggacagcctg ttttccgtgg gcgtattcag ggagccaata ttaatgacca	360
agcaaatact ggaattgacg ggcttgacag ttggcgagtt gccagctctc aagaaacgct	420
aaatgtccct gtcacaacct ttggttaaact gaccctgcc gcaggtactt tcaactgcgac	480
cttctacgtt cagcagtatc aaaac	505

<210> 2

<211> 165

<212> PRT

<213> Salmonella enteritidis

<400> 2

Met	Arg	Lys	Ser	Ala	Ser	Ala	Val	Ala	Val	Leu	Ala	Leu	Ile	Ala	Cys
1				5					10					15	

Gly Ser Ala His Ala Ala Gly Phe Val Gly Asn Lys Ala Glu Val Gln

20 25 30

Ala Ala Val Thr Ile Ala Ala Gln Asn Thr Thr Ser Ala Asn Trp Ser  
35 40 45

Gln Asp Pro Gly Phe Thr Gly Pro Ala Val Ala Gly Gln Lys Val  
50 55 60

Gly Thr Leu Ser Ile Thr Ala Thr Gly Pro His Asn Ser Val Ser Ile  
65 70 75 80

Ala Gly Lys Gly Ala Ser Val Ser Gly Gly Val Ala Thr Val Pro Phe  
85 90 95

Val Asp Gly Gln Gly Gln Pro Val Phe Arg Gly Arg Ile Gln Gly Ala  
100 105 110

Asn Ile Asn Asp Gln Ala Asn Thr Gly Ile Asp Gly Leu Ala Gly Trp  
115 120 125

Arg Val Ala Ser Ser Gln Glu Thr Leu Asn Val Pro Val Thr Thr Phe  
130 135 140

Gly Lys Ser Thr Leu Pro Ala Gly Thr Phe Thr Ala Thr Phe Tyr Val  
145 150 155 160

Gln Gln Tyr Gln Asn  
165

<210> 3  
<211> 128  
<212> PRT  
<213> Salmonella enteritidis

<400> 3

Ala Ala Gln Asn Thr Thr Ser Ala Asn Trp Ser Gln Asp Pro Gly Phe  
1 5 10 15

Thr Gly Pro Ala Val Ala Ala Gly Gln Lys Val Gly Thr Leu Ser Ile  
20 25 30

Thr Ala Thr Gly Pro His Asn Ser Val Ser Ile Ala Gly Lys Gly Ala

35 40 45

Ser Val Ser Gly Gly Val Ala Thr Val Pro Phe Val Asp Gly Gln Gly  
50 55 60

Gln Pro Val Phe Arg Gly Arg Ile Gln Gly Ala Asn Ile Asn Asp Gln  
65 70 75 80

Ala Asn Thr Gly Ile Asp Gly Leu Ala Gly Trp Arg Val Ala Ser Ser  
85 90 95

Gln Glu Thr Leu Asn Val Pro Val Thr Thr Phe Gly Lys Ser Thr Leu  
100 105 110

Pro Ala Gly Thr Phe Thr Ala Thr Phe Tyr Val Gln Gln Tyr Gln Asn  
115 120 125

<210> 4  
<211> 493  
<212> PRT  
<213> Salmonella enteritidis

<400> 4

Leu Thr Gln Asn Asn Leu Asn Lys Ser Gln Ser Ser Leu Ser Ser Ala  
1 5 10 15

Ile Glu Arg Leu Ser Ser Gly Leu Arg Ile Asn Ser Ala Lys Asp Asp  
20 25 30

Ala Ala Gly Gln Ala Ile Ala Asn Arg Phe Thr Ser Asn Ile Lys Gly  
35 40 45

Leu Thr Gln Ala Ser Arg Asn Ala Asn Asp Gly Ile Ser Ile Ala Gln  
50 55 60

Thr Thr Glu Gly Ala Leu Asn Glu Ile Asn Asn Asn Leu Gln Arg Val  
65 70 75 80

Arg Glu Leu Ser Val Gln Ala Thr Asn Gly Thr Asn Ser Asp Ser Asp  
85 90 95

Leu Lys Ser Ile Gln Asp Glu Ile Gln Gln Arg Leu Glu Glu Ile Asp

Arg	Val	Ser	Asn	Gln	Thr	Gln	Phe	Asn	Gly	Val	Lys	Val	Leu	Ser	Gln
		115					120					125			
Asp	Asn	Gln	Met	Lys	Ile	Gln	Val	Gly	Ala	Asn	Asp	Gly	Glu	Thr	Ile
	130					135					140				
Thr	Ile	Asp	Leu	Gln	Lys	Ile	Asp	Val	Lys	Ser	Leu	Gly	Leu	Asp	Gly
145					150					155					160
Phe	Asn	Val	Asn	Gly	Pro	Lys	Glu	Ala	Thr	Val	Gly	Asp	Leu	Lys	Ser
				165					170					175	
Ser	Phe	Lys	Asn	Val	Thr	Gly	Tyr	Asp	Thr	Tyr	Ala	Ala	Gly	Ala	Asp
			180					185					190		
Lys	Tyr	Arg	Val	Asp	Ile	Asn	Ser	Gly	Ala	Val	Val	Thr	Asp	Ala	Ala
		195					200					205			
Ala	Pro	Asp	Lys	Val	Tyr	Val	Asn	Ala	Ala	Asn	Gly	Gln	Leu	Thr	Thr
	210					215					220				
Asp	Asp	Ala	Glu	Asn	Asn	Thr	Ala	Val	Asp	Leu	Phe	Lys	Thr	Thr	Lys
225					230					235					240
Ser	Thr	Ala	Gly	Thr	Ala	Glu	Ala	Lys	Ala	Ile	Arg	Gly	Ala	Ile	Lys
				245					250					255	
Gly	Gly	Lys	Glu	Gly	Asp	Thr	Phe	Asp	Tyr	Lys	Gly	Val	Thr	Phe	Thr
			260					265					270		
Ile	Asp	Thr	Lys	Thr	Gly	Asp	Asp	Gly	Asn	Gly	Lys	Val	Ser	Thr	Thr
		275					280					285			
Ile	Asn	Gly	Glu	Lys	Val	Thr	Leu	Thr	Val	Ala	Asp	Ile	Ala	Thr	Gly
	290					295					300				
Ala	Thr	Asp	Val	Asn	Ala	Ala	Thr	Leu	Gln	Ser	Ser	Lys	Asn	Val	Tyr
305					310					315					320







1                    5                    10                    15

Glu Lys Val Thr Leu Thr Val Ala Asp Ile Ala Thr Gly Ala Thr Asp  
20                    25                    30

Val Asn Ala Ala Thr Leu Gln Ser  
35 40

```
<210>      8
<211>     27
<212>     PRT
<213>     Salmonella enteritidis
```

<400> 8

Asp Gly Asn Gly Lys Val Ser Thr Thr Ile Asn Gly Glu Lys Val Thr  
1 5 10 15

Leu Thr Val Ala Asp Ile Ala Thr Gly Ala Thr  
20 25

```
<210> 9
<211> 11
<212> PRT
<213> Salmonella enteritidis
```

<400> 9

Ser Thr Thr Ile Asn Gly Glu Lys Val Thr Leu  
1 5 10

<210>	10
<211>	22
<212>	DNA
<213>	Artificial Sequence

```
<220>
<223> Forward Primer for the C128 fragment of the fimbrial SEF14 antigen
```

```
<400> 10
tgcagctcag aatacaacat ca 22
```

```
<210> 11
<211> 21
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Reverse Primer for the C128 fragment of the fimbrial SEF14 antigen
```

```
<400> 11
gttttgatac tgctgaacgt a 21
```

<210>	12
<211>	26
<212>	DNA
<213>	Artificial Sequence

<220>  
<223> Forward Primer

```
<400> 12
aaggatccaa aggtggttaag gaagga                26
```

```
<210> 13
<211> 26
<212> DNA
<213> Artificial Sequence
```

<220>  
<223> Reverse Primer

```
<400> 13
ccgaattctt tgtcaccggt cactac 26
```

<210>	14
<211>	26
<212>	DNA
<213>	Artificial Sequence

<220>  
<223> Forward Primer

```
<400> 14
aaggatccaa aactggtgat gacggt 26
```

<210>	15
<211>	26
<212>	DNA
<213>	Artificial Sequence

<220>  
<223> Reverse Primer

```
<400> 15
ccgaattcgc ttgattgtaa ggtagc 26
```

<210> 16  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Forward Primer

<400> 16  
 aaggatccga cggtaatggt aaggtt

26

<210> 17  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Reverse Primer

<400> 17  
 ccgaattccg tcgcgccagt ggcaat

26

<210> 18  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Forward Primer

<400> 18  
 aaggatcctc tactaccatc aatggt

26

<210> 19  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Reverse Primer

<400> 19  
 ccgaattcta acgtaacttt ttcacc

26